

Title	Coprophagy in wild bonobos (<i>Pan paniscus</i>) at Wamba in the Democratic Republic of the Congo: a possibly adaptive strategy?
Author(s)	Sakamaki, Tetsuya
Citation	Primates (2010), 51(1): 87-90
Issue Date	2010-01
URL	http://hdl.handle.net/2433/123417
Right	The original publication is available at www.springerlink.com
Type	Journal Article
Textversion	publisher

1 Coprophagy in wild bonobos (*Pan paniscus*) at Wamba in the Democratic Republic of
2 the Congo: a possibly adaptive strategy?

3

4 Author's name: Tetsuya Sakamaki

5

6 Affiliation of author: Primate Research Institute, Kyoto University

7 Address of author: Primate Research Institute, Kyoto University, Kanrin 41, Inuyama,

8 Aichi 484-8506, Japan

9 E-mail: sakamaki@pri.kyoto-u.ac.jp

10 Tel: +81-568-63-0541

11 Fax: +81-568-63-0565

12

13

14 **Abstract** Four cases of coprophagy and two cases of fecal inspection were
15 identified during the 1,142 h of observing wild bonobos at Wamba in the Luo
16 Scientific Reserve in the Democratic Republic of the Congo. At least 5 females in the
17 study group practiced coprophagy and/or fecal inspection. According to our daily
18 behavioral observations, boredom and stress, insufficient roughage, and the search for
19 essential nutrients could not explain the coprophagy. Several episodes observed in this
20 study indicated that bonobos might have sought and ingested certain valuable food
21 items, such as hard *Dialium* seeds, in feces during relatively lean seasons. Although
22 coprophagy occurred only rarely among wild bonobos, this practice appeared to
23 represent a possibly adaptive feeding strategy during periods of food scarcity rather
24 than a behavioral abnormality.

25

26 **Keywords** Bonobo, *Pan paniscus*, Coprophagy, Inspect feces, Adaptive strategy,
27 Wamba.

28

29 **Introduction**

30

31 Coprophagy refers to the ingestion of one's own (autocoprophagy) or others'
32 (allocoprophagy) fecal material; this practice occurs in wild populations of lagomorphs,
33 rodents, and, to a lesser degree, piglets, foals, dogs, and primates (Soave and Brand
34 1991; Fish et al. 2007). African apes, gorillas, and chimpanzees practice coprophagy
35 both in captivity and in their natural environments (Harcourt and Stewart 1978; Akers
36 and Schildkraut 1985; Goodall 1986; Hook et al. 2002). Although coprophagy in wild
37 chimpanzees has been seen only rarely, this practice has been observed in several
38 long-term study sites, including those in Gombe, Mahale, Assirik, Fongoli, and Semliki
39 (Nishida et al. 1999; Payne et al. 2008).

40 Although the cause of coprophagy in chimpanzees and gorillas remains unclear,
41 several hypotheses have been proposed. (1) Boredom and stress: In captivity, fewer
42 social stimuli and less time to search for food can cause boredom and stress. Such
43 situations might induce coprophagy or other abnormal behaviors (Akers and
44 Schildkraut 1985; Nash et al. 1999). It has been hypothesized that coprophagy among
45 wild mountain gorillas might be associated with periods of heavy rain because gorillas
46 might need to relieve boredom and/or eat something warm under these circumstances
47 (Harcourt and Stewart 1978). (2) Insufficient roughage: Deficiencies in sources of
48 dietary roughage (e.g., fibrous leaves) can occur in captivity and might increase the
49 frequency of coprophagy (Fritz et al. 1992). (3) Essential nutrients: In some
50 lagomorphs and rodents, coprophagy is apparently an adaptive trait permitting full
51 digestion of complex carbohydrates. It also provide vitamins, minerals, amino acids,
52 and other nutrients (Soave and Brand 1991). Vitamin B12 might be a key reason for
53 engaging in coprophagy among primarily herbivorous apes because it is present only in
54 animal matter (Oxnard 1966). (4) Food scarcity: Chimpanzees at Gombe in Tanzania
55 engaged in coprophagy during the 1981 dry season, a period of fruit scarcity (Goodall
56 1986). The pressures of foraging in food-scarce environments might induce
57 coprophagy. (5) Reingesting valuable food items such as meat and hard seeds: Hard
58 seeds such as *Dialium* spp. and *Saba comorensis* might represent the target in the
59 reingested feces; that is, coprophagy might constitute a potentially efficient way of
60 ingesting the nutritional content of seeds that have already passed through the stomach
61 (Uehara 1979; Goodall 1986; Krief et al. 2004; Payne et al. 2008).

62 Although coprophagy has been observed among wild bonobos (*Pan paniscus*) at
63 Wamba and Lomako (ethogram created in a workshop, “Behavior, Ecology and
64 Conservation of Wild Bonobos: Current Activities and Plans for the Future,” Inuyama,
65 Japan, 2003), this is the first report to include details about coprophagy among wild

66 bonobos. I examined the applicability of the aforementioned hypotheses to my
67 observations.

68

69 **Methods**

70

71 I studied bonobos in the E1 group at Wamba (0°11'8"N, 22°37'58"E) in the northern
72 sector of the Luo Reserve in the Democratic Republic of the Congo. All individuals in
73 the group were identified and well habituated. Artificial provisioning was abolished in
74 1996. The history of the E1 group and the details of the study site have been described
75 by Kano (1992), Furuichi et al. (1998), Hashimoto et al. (2008) and Idani et al. (2008).

76 Observations were made during three study periods. Period 1: August
77 11–November 2, 2007; Period 2: January 31–March 11, 2008; Period 3: September 1,
78 2008–January 4, 2009. I attempted to locate the E1 group during 6 days of each week
79 and to follow the parties from one sleeping site to the next. I recorded *ad libitum* the
80 behaviors of those bonobos within sight as I followed them. Total observation time was
81 1,141 h 57 min across 156 days (Table 1). The E1 group was comprised of 23–26
82 individuals including 9 adult males and 6–7 adult females (15 years or older) during
83 the period under investigation.

84 In this study, coprophagy refers to feces-eating behavior, irrespective of
85 whether bonobos ate the feces or extracted and ate something selectively, or object of
86 ingestion was not confirmed.

87

88 **Results**

89

90 I observed four episodes of autocoprophagy (cases 1–4) and two episodes of fecal
91 inspection (cases^a 1 and 2) (Table 1). Coprophagy occurred 0.35 times per 100 hours.

92 Four females (2 adults, 1 subadult, and 1 immature bonobo) engaged autocoprophagy.
93 During the episodes of fecal inspection, 2 adult females (one of whom, Sala, practiced
94 coprophagy as well) defecated directly into their own hands, placed the feces close to
95 their faces, and then dropped it without ingesting.

96 Cases 1 and 2 occurred in October, a month of high fruit availability, including
97 fruits of *Landolphia* spp. (Apocynaceae), a favorite of bonobos. On the other hand,
98 cases 3 and 4 occurred in December, a month in which fruit was relatively less
99 plentiful (Kano and Mulavwa 1984; Mulavwa et al. 2008). When cases 3 and 4
100 occurred, most individuals in the E1 group showed symptoms of a flu-like disease (i.e.,
101 coughing, sneezing, and/or nose-picking) (Sakamaki et al. 2009). During this period of
102 time, the E1 group members divided into relatively small parties and did not travel
103 long distances each day. They ate a relatively restricted diet, which consisted primarily
104 of fruits of *Dialium pachyphyllum*, *Dialium zenkeri* (Caesalpiniaceae), and *Musanga*
105 *cecropioides* (Moraceae).

106 All feeding behaviors involved in coprophagy occurred high in the trees,
107 rendering it difficult to observe the behavioral patterns in detail. However, case 4 of
108 coprophagy was clearly observed; in this instance, Fuku, a subadult female,
109 confidently used her lips to extract *Dialium* seeds from the feces in her hand, ate the
110 seeds, and discarded other fibrous parts in the feces. In case^a 2 and Case 4, bonobos
111 held their feces in their hands to inspect and/or consume it. After dropping the feces,
112 they rubbed their hands on the stems of a tree/vine. I did not see diarrheic individuals
113 inspect and/or eat their feces, and most feces involved in the episodes seemed to be
114 hard.

115

116 **Discussion**

117

118 This is the first report on the details of coprophagy among wild bonobos. This study
119 indicated that bonobos only rarely engaged in coprophagy in natural environments.
120 Although we cannot infer the cause of coprophagy on the basis of the data collected in
121 this study, we can rule out certain previous causal hypotheses. Hypothesis 1, boredom
122 and stress: We observed the subjects in their natural habitat, which involved searching
123 for foods and interacting with group members. This environment was no less socially
124 stimulating than is captivity. In addition, boredom during heavy rains could not explain
125 the coprophagy observed in this study because coprophagy occurred outside of the
126 months with the greatest rainfall (September-November; Mulavwa et al. 2008). Being
127 followed by human observers almost all day might have been stressful to the bonobos.
128 However, the bonobos in the E1 group were so accustomed to humans that they did not
129 seem to be unduly stressed by being followed by the observers. Hypothesis 2, lack of
130 dietary roughage: This hypothesis could not explain the coprophagy observed in this
131 study because sources of roughage were almost always available in the habitat; for
132 example, piths/shoots of terrestrial herb vegetation [e.g., *Aframomum* spp.
133 (*Zingiberaceae*), *Haumania Liebrechtsiana* and *Megaphrynium macrostachyum*
134 (*Marantaceae*)] and young leaves [e.g., *Scorodophloeus zenkeri* (*Caesalpinaceae*)]
135 were observed (Kano and Mulavwa 1984). Hypothesis 3, essential nutrients: Bonobos
136 in the study group were able to consume essential nutrients such as Vitamin B12 by
137 eating insect larvae, earthworms, vertebrates, and eggs of birds (Kano and Mulavwa
138 1984). Even if certain nutrients were derived from feces, the concentration of these
139 nutrients in the feces would be insufficient for daily requirements. Further studies are
140 needed to investigate nutrients in foods and feces.

141 The other hypotheses suggest variables that might have induced coprophagy
142 among the bonobos in this study. Hypothesis 4, food scarcity: Contrary to expectation
143 of this hypothesis, coprophagy observed in cases 1 and 2 occurred when favorite fruit

144 foods were relatively abundant. However, coprophagy in cases 3 and 4 occurred when
145 fruit was less plentiful and when the bonobos were traveling only a short distance, and
146 therefore, feeding activities were largely limited due to the epidemic of a flu-like
147 disease. Hypothesis 5, reingesting valuable food items such as hard seeds: Rogers et al.
148 (1998) reported that wild gorillas extract and consume *Dialium* seeds from feces, and
149 Krief et al. (2004) also reported that chimpanzees rehabilitated into a natural
150 environment ingested *Dialium* seeds extracted from feces. Certain hard seeds in feces,
151 such as *Dialium* spp., might be items of interest for gorillas and chimpanzees because
152 of their protein content (Krief et al. 2004; Payne et al. 2008). During the season that
153 included coprophagy, cases 1 and 2 and fecal inspection cases^a 1 and 2, bonobos
154 primarily ate fruits of the *Landolphia* spp., the seeds of which are similar to those of
155 *Saba comorensis* that chimpanzees crunch and eat (Uehara 1979; Payne et al. 2008).
156 During the season including cases 3 and 4, bonobos primarily ate fruits of *Dialium* spp.,
157 and a bonobo was observed to extract the seeds in case 4. In case^a 2 and case 4,
158 bonobos rubbed their hands on the stems of a tree/vine probably to clean their hands
159 after they dropped the feces. These observations suggested that bonobos searched for
160 certain valuable food items in their feces during relatively lean food seasons.

161

162 **Acknowledgements** This study was financially supported by the Global
163 Environment Research Fund (F-061 to T. Nishida) of the Japanese Ministry of the
164 Environment. I thank the Research Center for Ecology and Forestry (CREF) of the
165 Democratic Republic of the Congo for permission to conduct the fieldwork in the Luo
166 Scientific Reserve. I am also grateful to N. Mwanza, M. Mulavwa, K. Yangozene, S.
167 Kuroda, H. Takemoto, Y. Tsuji, the research assistants at Wamba, and other members
168 of CREF for their support and cooperation in the field, as well as to T. Nishida, T.
169 Furuichi, and members of the Wamba Committee for Bonobo Research for their

170 encouragement and useful comments.

171

172 **References**

173

174 Akers JS, Schildkraut DS (1985) Regurgitation/reingestion and coprophagy in captive
175 gorillas. *Zoo Biol* 4: 99-109

176 Fish KD, Sauther ML, Loudon JE, Cuzzo FP (2007) Coprophagy by wild ring-tailed
177 lemurs (*Lemur catta*) in human-disturbed locations adjacent to the Beza
178 Mahafaly Special Reserve, Madagascar. *Am J Primatol* 69:713-718

179 Fritz J, Maki S, Nash LT, Martin T, Matevia M (1992) The relationship between forage
180 material and levels of coprophagy in captive chimpanzees (*Pan troglodytes*). *Zoo*
181 *Biol* 11: 313-318

182 Furuichi T, Idani G, Ihobe H, Kuroda S, Kitamura K, Mori A, Enomoto T, Okayasu N,
183 Hashimoto C, Kano T (1998) Population dynamics of wild bonobos (*Pan*
184 *paniscus*) at Wamba. *Int J Primatol* 19: 1029-1043

185 Goodall J (1986) The chimpanzees of Gombe: Patterns of behavior. Harvard
186 University Press, Cambridge.

187 Harcourt AH, Stewart KJ (1978) Coprophagy by wild mountain gorilla. *East Afr Wildl*
188 *J* 16: 223-225

189 Hashimoto C, Tashiro Y, Hibino E, Mulavwa M, Yangozene K, Furuichi T, Idani G,
190 Takenaka O (2008) Longitudinal structure of a unit-group of bonobos: Male
191 philopatry and possible fusion of unit-groups. In: Furuichi T, Thompson J (eds.)
192 The Bonobos: Behavior, ecology, and conservation. Springer, New York, pp
193 107-119

194 Hook MA, Lambeth SP, Perlman JE, Stavisky R, Bloomsmith MA, Schapiro SJ (2002)
195 Inter-group variation in abnormal behavior in chimpanzees (*Pan troglodytes*) and

196 rhesus macaques (*Macaca mulatta*). *Appl Anim Behav Sci* 76: 165-176

197 Idani G, Mwanza N, Ihobe H, Hashimoto C, Tashiro Y, Furuichi T (2008) Changes in
198 the status of bonobos, their habitat, and the situation of humans at Wamba in the
199 Luo Scientific Reserve, Democratic Republic of Congo. In: Furuichi T,
200 Thompson J (eds.) *The Bonobos: Behavior, ecology, and conservation*. Springer,
201 New York, pp 291-302

202 Kano T (1992) *The Last Ape: Pigmy Chimpanzee Behavior and Ecology*. Stanford
203 Univ Press, Stanford.

204 Kano T, Mulavwa M (1984) Feeding ecology of the pygmy chimpanzees (*Pan*
205 *paniscus*) of Wamba. In: Susman RL (ed) *The Pygmy Chimpanzee*. Plenum
206 Publishing Corporation, pp. 233-274

207 Krief S, Jamart A, Hladik CM (2004) On the possible adaptive value of coprophagy in
208 free-ranging chimpanzees. *Primates* 45: 141-145

209 Mulavwa M, Furuichi T, Yangozene K, Yamba-Yamba M, Motema-Salo B, Idani G,
210 Ihobe H, Hashimoto C, Tashiro Y, Mwanza N (2008) Seasonal change in fruit
211 production and party size of bonobos at Wamba. In: Furuichi T, Thompson J
212 (eds.) *The Bonobos: Behavior, ecology, and conservation*. Springer, New York,
213 pp 121-134

214 Nash LT, Fritz J, Alford PA, Brent L (1999) Variables influencing the origins of diverse
215 abnormal behaviors in a large sample of captive chimpanzees (*Pan troglodytes*).
216 *Am J Primatol* 48: 15-29

217 Nishida T, Kano T, Goodall J, McGrew WC, Nakamura M (1999) Ethogram and
218 ethnography of Mahale chimpanzees. *Anthropol Sci* 107: 141-188

219 Oxnard CE (1966) Vitamin B12 nutrition in some primates in captivity. *Folia Primatol*
220 4: 424-431

221 Payne CLR, Webster TH, Hunt KD (2008) Coprophagy by the semi-habituated

222 chimpanzees of Semliki, Uganda. *Pan Afr News* 15: 29-32

223 Rogers ME, Voysey BC, McDonald KE, Parnell RJ, Tutin CEG (1998) Lowland
224 gorillas and seed dispersal: The importance of nest sites. *Am J Primatol* 45:
225 45-68

226 Sakamaki T, Mulavwa M, Furuichi T (2009) Flu-like Epidemics in Wild Bonobos (*Pan*
227 *paniscus*) at Wamba, the Luo Scientific Reserve, Democratic Republic of Congo.
228 *Pan Afr News* 16: 1-4

229 Soave O, Brand CD (1991) Coprophagy in animals: A review. *Cornell Vet* 81: 357-64

230 Uehara S (1979) The chimpanzees of Kasoge K group [in Japanese]. *Monkey* 170:
231 16-25

Table 1. Observations of coprophagy

Month and Year	Observation		Observation of coprophagy		
	No. days	Time			
Period 1					
August 2007	10	61 h 57 min	None	-	-
September 2007	21	137 h 9 min	None	-	-
October 2007	22	157 h 10 min	Case 1 (Oct 31)	Sala (adult female)	At 0921 h, she took her feces in her left hand and ate it. She again took her feces in her left hand and ate it. The feces seemed to be dark in color and hard.
November 2007	2	13 h 10 min	None	-	-
Period 2					
January 2008	1	10 h 58 min	None	-	-
February 2008	17	119 h 47 min	None	-	-
March 2008	7	66 h 58 min	None	-	-
Period 3					
September 2008	22	151 h 7 min	Case ^a 1 (Sep 24)	Sala (adult female)	At 0821 h, she evacuated a small amount of feces, which fell on the ground. She held the fecal material that emerged second in her left hand, moved it close to her face, looked at it carefully, and then dropped it.
October 2008	19	148 h 6 min	Case ^a 2 (Oct 10)	Hoshi (adult female)	At 1003 h, she held her feces in her right hand after appearing to experience some difficulty with the excretory process, moved it close to her face and looked at it, and then dropped it. Next, she rubbed her right hand on a woody vine.
			Case 2 (Oct 10)	Jacky (adult female)	At 1439 h, her feces fell between her fingers even though she positioned her hand close to her anus. She held the second feces in her left hand, moved it close to her face, looked at it, and then ate it.
November 2008	8	59 h 16 min	None	-	-
December 2008	25	211 h 43 min	Case 3 (Dec 5)	Nachi (2-year-old female)	At 0620 h, she grasped her feces in her right hand and ate it. The feces were about 5 cm long and seemed to be light in color.
			Case 4 (Dec 10)	Fuku (approximately 10-year-old female)	At 0731 h, she grasped her feces in her left hand. At 0732 h, she moved it close to her face and looked at it. At 0735 h, she ate it and dropped something that resembled seeds from her mouth. At 0736 h, she rubbed the palm of her left hand on a tree trunk. At 0741 h, she kept the feces in her hand while crunching seeds in her mouth. She used her lips to select only seeds from the feces, and chewed them in her mouth. She dropped other fibers or material from the feces. At 0747 h, she finished eating the feces and rubbed her left hand on a tree trunk. The feces seemed to be ochreous in color.
January 2009	2	4 h 36 min	None	-	-
Total	156	1141 h 57 min			